# UNCLASSIFIED

AD 268 331

Reproduced by the

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U.S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

\(\langle 2 - 1 - 2 \)
\(\text{XEROX}\)

XEROX

ZPh-120 Physics Section

CATALOGED BY ASTIK 268331 AS AD NO.

INFRARED EMISSIVITY OF OH

W. Malkmus

September 1961

268 331



# INFRARED EMISSIVITY OF OH

W. Malkmus

September 1961

This work was conducted under Geophysics Research Directorate Air Force Cambridge Research Laboratories Contract No. AF 19 (604)-5554 ARPA Order No. 116, Project Defender



## INFRARED EMISSIVITY OF OH

### W. Malkmus

In a previous publication, equations were developed for the computation of diatomic gas emissivities for an assumed model of an anharmonic vibrating rotator with the first approximation to the vibration-rotation interaction. This paper is an application of the results of that analysis to the OH molecule.

The fundamental band strength of the OH molecule was taken<sup>2</sup> as 100 cm<sup>-2</sup> atm<sup>-1</sup> at 300 K. In the absence of experimental data, the factor 0 (the ratio of the zeroth to the first order term in the expansion of the electric dipole moment) and the average line width (which enters only in the strong line approximation) were assumed to be the same as for the HF molecule.

The results are presented in Figures 1 to 16, for both the weak line and the strong line approximations. Computations have also been made for the case of pure Doppler broadening and will be published in a separate report along with such computations for other diatomic molecules.<sup>3</sup>

### REFERENCES

- 1. W. Malkmus and A. Thomson, Convair Report ZPh-095, May 5, 1961; also in Journal of Molecular Spectroscopy and Radiative Transfer (to be published).
- 2. W. S. Benedict and E. K. Plyler, "High-Resolution Spectra of Hydrocarbon Flames," in <u>Energy Transfer in Hot Gases</u>, pp. 57-73, N. B. S. Circular No. 523, Washington, D. C., 1954.
- 3. W. Malkmus and A. Thomson, Convair Report ZPh-119 (to be published).




FIGURE 1. SPECTRAL EMISSIVITY OF OH AT T = 300°K (Weak Line Approximation)




FIGURE : SPECTRAL EMISSIVITY OF OH AT T = 600 K (Weak Line Approximation)




FIGURE 3. SPECTRAL EMISSIVITY OF OH AT T = 1200°K (Weak Line Approximation)




FIGURE 4. SPECTRAL EMISSIVITY OF OH AT T = 1800°K (Weak Line Approximation)




FIGURE 5. SPECTRAL EMISSIVITY OF OH AT T = 2400 °K (Weak Line Approximation)




FIGURE 6. SPECTRAL EMISSIVITY OF OH AT T = 3000 °K (Weak Line Approximation)




FIGURE 7. SPECTRAL EMISSIVITY OF OH AT T = 5000°K (Weak Line Approximation)




FIGURE 8. SPECTRAL EMISSIVITY OF OH AT T = 7000°K (Weak Line Approximation)




FIGURE 9. SPECTRAL EMISSIVITY OF OH AT T = 300°K (Strong Line Approximation)




FIGURE 10. SPECTRAL EMISSIVITY OF OH AT T = 600°K (Strong Line Approximation)




FIGURE 11. SPECTRAL EMISSIVITY OF OH AT T = 1200°K (Strong Line Approximation)




FIGURE 12. SPECTRAL EMISSIVITY OF OH AT T = 1800°K (Strong Line Approximation)





FIGURE 13. SPECTRAL EMISSIVITY OF OH AT T = 2400 °K (Strong Line Approximation)



8




FIGURE 15. SPECTRAL EMISSIVITY OF OH AT T = 5000°K (Strong Line Approximation)




FIGURE 16. SPECTRAL EMISSIVITY OF OH AT T = 7000°K (Strong Line Approximation)